



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A2	(11) International Publication Number: WO 98/50584 (43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/08926 (22) International Filing Date: 1 May 1998 (01.05.98) (30) Priority Data: 60/045,400 2 May 1997 (02.05.97) US (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA as represented by THE SECRETARY OF THE DEPARTMENT OF HEALTH AND HUMAN SERVICES, c/o Centers for Disease Control and Prevention, Technology Transfer Office [US/US]; Atlanta, GA 30329 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MORRISON, Christine, J. [US/US]; 3110 Tolbert Drive, Decatur, GA 30033 (US). REISS, Errol [US/US]; 3642 Castaway Court, Chamblee, GA 30341 (US). AIDOREVICH, Liliana [VE/VE]; Calle Circunvalación, Manzana B7 #16 Urlo, El Castan Maracay Edo Aragua (VE). CHOI, Jong, Soo [KR/KR]; 202-1506, Sinchungi-Town Apartment, Whangum-dong, Susung-gu, Taegu City 706-040 (KR). (74) Agents: WARREN, William, L. et al.; Jones & Askew, 37th floor, 191 Peachtree Street, N.E., Atlanta, GA 30303 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NUCLEIC ACIDS FOR DETECTING <i>ASPERGILLUS</i> SPECIES AND OTHER FILAMENTOUS FUNGI (57) Abstract <p>Nucleic acids for detecting <i>Aspergillus</i> species and other filamentous fungi are provided. Unique internal transcribed spacer 2 coding regions permit the development of nucleic acid probes specific for five different species of <i>Aspergillus</i>, three species of <i>Fusarium</i>, four species of <i>Mucor</i>, two species of <i>Penicillium</i>, five species of <i>Rhizopus</i>, one species of <i>Rhizomucor</i>, as well as probes for <i>Absidia corymbifera</i>, <i>Cunninghamella elagans</i>, <i>Pseudallescheria boydii</i>, and <i>Sporothrix schenckii</i>. The invention thereby provides methods for the species-specific detection and diagnosis of infection by <i>Aspergillus</i>, <i>Fusarium</i>, <i>Mucor</i>, <i>Penicillium</i>, <i>Rhizopus</i>, <i>Rhizomucor</i>, <i>Absidia</i>, <i>Cunninghamella</i>, <i>Pseudallescheria</i> or <i>Sporothrix</i> in a subject. Furthermore, genus-specific probes are also provided for <i>Aspergillus</i>, <i>Fusarium</i> and <i>Mucor</i>, in addition to an all-fungus nucleic acid probe.</p>		

Best Available Copy

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

10

**NUCLEIC ACIDS FOR DETECTING
ASPERGILLUS SPECIES AND OTHER FILAMENTOUS FUNGI**

15 This invention was made in the Centers for Disease Control
Mycotic Diseases Laboratories, an agency of the United States Government.

Technical Field

20 This application relates in general to the field of diagnostic
microbiology. In particular, the invention relates to the species-specific
detection of *Aspergillus*, *Fusarium*, *Mucor*, *Penicillium*, *Rhizopus*,
Rhizomucor, *Absidia*, *Cunninghamella*, *Pseudallescheria boydii*
(*Scedosporium apiospermum*), and *Sporothrix* species.

Background of the Invention

25 In recent years, chemotherapy for hematological
malignancies, and high-dose corticosteroid treatment for organ transplant
recipients, along with the spread of AIDS, have greatly increased the number
of immunocompromised patients (1, 12, 14, 43). Saprophytic filamentous
fungi, such as *Aspergillus*, *Rhizopus*, and *Mucor* species, found in the
30 environment and considered to be of low virulence, are now responsible for
an increasing number of infections in the immunocompromised host (17, 20,
43). In addition, these infections are often fulminant and rapidly fatal in
immunocompromised patients (7, 11, 12, 20, 44). Morbidity and mortality
is extremely high; for example, aspergillosis has a mortality rate of
35 approximately 90% (8, 11).

To complicate matters, diagnosis is difficult and symptoms
are often non-specific (18, 27, 29, 42, 44). Antibody-based tests can be

unreliable due to the depressed or variable immune responses of immunocompromised patients (2, 9, 18, 46). Antigen detection tests developed to date have fallen short of the desired sensitivity (2, 9, 38). Radiographic evidence can be non-specific and inconclusive (5, 29, 36), although some progress in diagnosis has been made with the advent of computerized tomography (40). However, definitive diagnosis still requires either a positive blood or tissue culture or histopathological confirmation (3, 21). An added complication is that the invasive procedures necessary to obtain biopsy materials are often not recommended in thrombocytopenic patient populations (37, 41).

Even when cultures of blood, lung or rhinocerebral tissues are positive, morphological and biochemical identification of filamentous fungi can require several days for adequate growth and sporulation to occur, delaying targeted drug therapy. Some atypical isolates may never sporulate, making identification even more difficult (23). When histopathology is performed on tissue biopsy sections, the morphological similarities of the various filamentous fungi in tissue make differentiation difficult (16). Fluorescent antibody staining of histopathological tissue sections is not specific unless cross-reactive epitopes are absorbed out which can make the resultant antibody reactions weak (14, 19). Therapeutic choices vary (7, 41, 44) making a test to rapidly and specifically identify filamentous fungi urgently needed for the implementation of appropriately targeted therapy. Early and accurate diagnosis and treatment can decrease morbidity and increase the chances for patient survival (6, 27, 39). Furthermore, identification of filamentous fungi to at least the species level would be epidemiologically useful (24, 31, 43, 47).

PCR-based methods of detection, which show promise as rapid, sensitive means to diagnose infections, have been used in the identification of DNA from *Candida* species (13, 15, 30) and some other fungi, particularly *Aspergillus* species (31, 33, 45). However, most of these tests are only genus-specific (28, 38) or are directed to detect only single-copy genes (4, 35). Others have designed probes to detect multi-copy genes so as to increase test sensitivity (31, 33) but in doing so have lost test specificity because they have used highly conserved genes, which detect one or a few species but which are also plagued with cross-reactivities to human, fungal or even viral DNA (25, 31, 33).

Therefore, it is an object of the invention to provide improved materials and methods for detecting and differentiating *Aspergillus* and other filamentous fungal species in the clinical and laboratory settings.

Summary of the Invention

The present invention relates to nucleic acids for detecting *Aspergillus*, *Fusarium*, *Mucor*, *Penicillium*, *Rhizopus*, *Rhizomucor*, *Absidia*, *Cunninghamella*, *Pseudallescheria* (*Scedosporium*), and *Sporothrix* species. Unique internal transcribed spacer 2 coding regions permit the development of probes specific for five different *Aspergillus* species, *A. flavus*, *A. fumigatus*, *A. niger*, *A. terreus*, and *A. nidulans*. The invention thereby provides methods for the species-specific detection and diagnosis of *Aspergillus* infection in a subject. In addition, species probes have been developed for three *Fusarium*, four *Mucor*, two *Penicillium*, five *Rhizopus* and one *Rhizomucor* species, as well as probes for *Absidia corymbifera*, *Cunninghamella elegans*, *Pseudallescheria boydii* (*Scedosporium apiospermum*), and *Sporothrix schenckii*. Generic probes for *Aspergillus*, *Fusarium*, and *Mucor* species have also been developed.

These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiments and the appended claims.

Detailed Description of the Invention

This invention provides a simple, rapid, and useful method for differentiating filamentous fungal species from each other and from other medically important fungi. This invention enables a rapid, simple and useful method to isolate fungal DNA from host samples, and to apply the species- and genus-specific probes for the diagnosis of a disease. Ultimately, these probes can be used for *in situ* hybridization or *in situ* PCR diagnostics so that the morphology of host tissue, and microorganisms, remain intact.

The invention provides nucleic acids containing regions of specificity for five *Aspergillus*, three *Fusarium*, four *Mucor*, two *Penicillium*, five *Rhizopus* and one *Rhizomucor* species as well as probes for *Absidia corymbifera*, *Cunninghamella elegans*, *Pseudallescheria boydii* (*Scedosporium apiospermum*), and *Sporothrix schenckii*. These nucleic acids are from the internal transcribed spacer 2 ("ITS2") region of ribosomal deoxyribonucleic

acid (rDNA) of the genome of the aforementioned filamentous fungi. The ITS2 region is located between the 5.8S rDNA region and the 28S rDNA region.

In particular, the invention provides nucleic acids from
5 *Aspergillus flavus* (SEQ ID NO:1), *Aspergillus fumigatus* (SEQ ID NO:2),
Aspergillus niger (SEQ ID NO:3), *Aspergillus terreus* (SEQ ID NO:4),
Aspergillus nidulans (SEQ ID NO:5), *Fusarium solani* (SEQ ID NO:6),
Fusarium moniliforme (SEQ ID NO:7), *Mucor rouxii* (SEQ ID NO:8), *Mucor*
10 *racemosus* (SEQ ID NO:9), *Mucor plumbeus* (SEQ ID NO:10), *Mucor indicus*
(SEQ ID NO:11), *Mucor circinilloides f. circinelloides* (SEQ ID NO:12),
Rhizopus oryzae (SEQ ID NO:13 and NO:14), *Rhizopus microsporus* (SEQ ID
NO:15 and 16), *Rhizopus circinans* (SEQ ID NO:17 and 18), *Rhizopus*
stolonifer (SEQ ID NO: 19), *Rhizomucor pusillus* (SEQ ID NO:20), *Absidia*
15 *corymbifera* (SEQ ID NO:21 and 22), *Cunninghamella elegans* (SEQ ID
NO:23), *Pseudallescheria boydii* (teleomorph of *Scedosporium apiospermum*)
(SEQ ID NO:24, 25, 26, and 27), *Penicillium notatum* (SEQ ID NO:28), and
Sporothrix schenkii (SEQ ID NO:29). These sequences can be used to identify
and distinguish the respective species of *Aspergillus*, *Fusarium*, *Mucor*,
20 *Rhizopus*, and *Penicillium*, and identify and distinguish these species from
each other and from *Absidia corymbifera*, *Cunninghamella elegans*,
Pseudallescheria boydii(*Scedosporium apiospermum*), and *Sporothrix*
schenkii.

Furthermore, the invention provides isolated nucleic acid
probes derived from GenBank nucleic acid sequences (for *Penicillium*
25 *marneffei* and *Fusarium oxysporum* only) or from the above nucleic acid
sequences which may be used as species-specific identifiers of *Aspergillus*
flavus (SEQ ID NO:30 and 31), *Aspergillus fumigatus* (SEQ ID NO:32),
Aspergillus niger (SEQ ID NO:33), *Aspergillus terreus* (SEQ ID NO:34),
Aspergillus nidulans (SEQ ID NO: 35), *Mucor rouxii* (SEQ ID NO:36),
30 *Mucor plumbeus* (SEQ ID NO:37), *Mucor indicus* (SEQ ID NO:38), *Mucor*
circinilloides f. circinelloides (SEQ ID NO:39), *Mucor racemosus* (SEQ ID
NO:40), *Rhizopus oryzae* (SEQ ID NO:41), *Rhizopus circinans* (SEQ ID
NO:42), *Rhizomucor pusillus* (SEQ ID NO:43), *Rhizopus stolonifer* (SEQ ID
NO:44), *Pseudallescheria boydii* (*Scedosporium apiospermum*)(SEQ ID
35 NO:45), *Penicillium notatum* (SEQ ID NO:46), *Penicillium marneffei* (SEQ
ID NO:47 and 48), *Fusarium moniliforme* (SEQ ID NO:49), *Fusarium*
oxysporum (SEQ ID NO:50), *Fusarium solani* (SEQ ID NO:51),

Cunninghamella elegans (SEQ ID NO: 52, 53, and 54), *Absidia corymbifera* (SEQ ID NO:55), *Sporothrix schenkii* (SEQ ID NO:56), and *Rhizopus microsporus* (SEQ ID NO:57). Such probes can be used to selectively hybridize with samples containing nucleic acids from species of *Aspergillus*, *Fusarium*, *Mucor*, *Rhizopus* (or *Rhizomucor*), *Penicillium*, or from *Absidia corymbifera*, *Cunninghamella elegans*, *Pseudallescheria boydii* (*Scedosporium apiospermum*), and *Sporothrix schenkii*. These fungi can be detected after polymerase chain reaction or ligase chain reaction amplification of fungal DNA and specific probing of amplified DNA with DNA probes labeled with digoxigenin, reacted with anti-digoxigenin antibodies labeled with horseradish peroxidase and a colorimetric substrate, for example. Additional probes can routinely be derived from the sequences given in SEQ ID NOs:1-29, which are specific for the respective species. Therefore, the probes shown in SEQ ID NOs:30-57 are only provided as examples of the species-specific probes that can be derived from SEQ ID NOs:1-29.

Generic probes for *Aspergillus* (SEQ ID NO:58), *Fusarium*, (SEQ ID NO:59) and *Mucor* (SEQ ID NO:60) species have also been developed to identify all members of their respective species which are listed above as well as an all-fungus biotinylated probe (SEQ ID NO:61) to capture all species-specific and generic probes listed above for their detection.

By "isolated" is meant nucleic acid free from at least some of the components with which it naturally occurs. By "selective" or "selectively" is meant a sequence which does not hybridize with other nucleic acids to prevent adequate determination of an *Aspergillus*, *Fusarium*, *Mucor*, *Penicillium*, *Rhizopus* or *Rhizomucor* genus or species or of *Absidia corymbifera*, *Cunninghamella elegans*, *Pseudallescheria boydii* (*Scedosporium apiospermum*), or *Sporothrix schenckii* species.

The hybridizing nucleic acid should have at least 70% complementarity with the segment of the nucleic acid to which it hybridizes. As used herein to describe nucleic acids, the term "selectively hybridizes" excludes the occasional randomly hybridizing nucleic acids and thus has the same meaning as "specifically hybridizing". The selectively hybridizing nucleic acids of the invention can have at least 70%, 80%, 85%, 90%, 95%, 97%, 98%, and 99% complementarity with the segment of the sequence to which it hybridizes.

The invention contemplates sequences, probes and primers which selectively hybridize to the complementary, or opposite, strand of DNA

as those specifically provided herein. Specific hybridization with nucleic acid can occur with minor modifications or substitutions in the nucleic acid, so long as functional species-specific or genus-specific hybridization capability is maintained. By "probe" is meant nucleic acid sequences that can be used as probes or primers for selective hybridization with complementary nucleic acid sequences for their detection or amplification, which probes can vary in length from about 5 to 100 nucleotides, or preferably from about 10 to 50 nucleotides, or most preferably about 18 nucleotides. The invention provides isolated nucleic acids that selectively hybridize with the species-specific nucleic acids under stringent conditions and should have at least 5 nucleotides complementary to the sequence of interest. *See generally, Maniatis (26).*

If used as primers, the invention provides compositions including at least two nucleic acids which hybridize with different regions so as to amplify a desired region. Depending on the length of the probe or primer, target region can range between 70% complementary bases and full complementarity and still hybridize under stringent conditions. For example, for the purpose of diagnosing the presence of the *Aspergillus*, the degree of complementarity between the hybridizing nucleic acid (probe or primer) and the sequence to which it hybridizes (e.g., *Aspergillus* DNA from a sample) is at least enough to distinguish hybridization with a nucleic acid from other yeasts and filamentous fungi. The invention provides examples of nucleic acids unique to each filamentous fungus in the listed sequences so that the degree of complementarity required to distinguish selectively hybridizing from nonselectively hybridizing nucleic acids under stringent conditions can be clearly determined for each nucleic acid.

Alternatively, the nucleic acid probes can be designed to have homology with nucleotide sequences present in more than one species of the fungi listed above. Such a nucleic acid probe can be used to selectively identify a group of species such as the generic probes listed for *Aspergillus* (SEQ ID NO:58), *Fusarium* (SEQ ID NO:59), and *Mucor* (SEQ ID NO:60) as well as all fungi listed (SEQ ID NO:61). Additionally, the invention provides that the nucleic acids can be used to differentiate the filamentous fungi listed in general from other filamentous fungi and yeasts, such as *Candida* species. Such a determination is clinically significant, since therapies for these infections differ.

The invention further provides methods of using the nucleic acids to detect and identify the presence of the filamentous fungi listed, or

particular species thereof. The method involves the steps of obtaining a sample suspected of containing filamentous fungi. The sample may be taken from an individual, such as blood, saliva, lung lavage fluids, vaginal mucosa, tissues, etc., or taken from the environment. The filamentous fungal cells can then be lysed, and the DNA extracted and precipitated. The DNA is preferably amplified using universal primers derived from the internal transcribed spacer regions, 18S, 5.8S and 28S regions of the filamentous fungal rDNA. Examples of such universal primers are shown below as ITS1 (SEQ ID NO: 62), ITS3 (SEQ ID NO: 63), ITS4 (SEQ ID NO: 64). Detection of filamentous fungal DNA is achieved by hybridizing the amplified DNA with a species-specific probe that selectively hybridizes with the DNA. Detection of hybridization is indicative of the presence of the particular genus (for generic probes) or species (for species probes) of filamentous fungus.

Preferably, detection of nucleic acid (e.g. probes or primers) hybridization can be facilitated by the use of detectable moieties. For example, the species-specific or generic probes can be labeled with digoxigenin, and an all-fungus probe, such as described in SEQ ID NO:61, can be labeled with biotin and used in a streptavidin-coated microtiter plate assay. Other detectable moieties include radioactive labeling, enzyme labeling, and fluorescent labeling, for example.

The invention further contemplates a kit containing one or more species-specific probes, which can be used for the detection of particular filamentous fungal species and genera in a sample. Such a kit can also contain the appropriate reagents for hybridizing the probe to the sample and detecting bound probe. The invention may be further demonstrated by the following non-limiting examples.

Examples

In this example, PCR assay employing universal, fungus-specific primers and a simple, rapid EIA-based format for amplicon detection were used.

Extraction of Filamentous Fungal DNA

A mechanical disruption method was used to obtain DNA from filamentous fungal species and an enzymatic disruption method described previously (13) was used to obtain DNA from yeasts. Filamentous fungi were grown for 4 to 5 days on Sabouraud dextrose agar slants (BBL,

division of Becton Dickinson, Cockeysville, MD) at 35°C. Two slants were then washed by vigorously pipeting 5 mls of 0.01 M potassium phosphate buffered saline (PBS) onto the surface of each slant and the washes were transferred to 500 ml Erlenmeyer flasks containing 250 ml of Sabouraud dextrose broth (BBL). Flasks were then incubated for 4 to 5 days on a rotary shaker (140 rpm) at ambient temperature. Growth was then harvested by vacuum filtration through a sterile Whatman #1 filter paper which had been placed into a sterile Buchner funnel attached to a 2 L side-arm flask. The resultant cellular mat was washed on the filtration apparatus three times with sterile distilled water, removed from the filter paper by gentle scraping with a rubber policeman, and placed into a sterile Petri plate which was then sealed with parafilm and frozen at -20°C until used.

Just prior to use, a portion of the frozen cellular mat, equal in size to a quarter, was removed and placed into a cold mortar (6" diameter). Liquid nitrogen was added to cover the mat which was then ground into a powder with a pestle. Additional liquid nitrogen was added as needed to keep the mat frozen during grinding.

DNA was then purified using proteinase K and RNase treatment, multiple phenol extractions, and ethanol precipitation by conventional means (26).

PCR amplification

The fungus-specific, universal primer pair ITS3 (5'-GCA TCG ATG AAG AAC GCA GC-3') (SEQ ID NO: 63) and ITS4 (5'-TCC TCC GCT TAT TGA TAT GC-3') (SEQ ID NO: 64) was used to amplify a portion of the 5.8S rDNA region, the entire ITS2 region, and a portion of the 28S rDNA region for each species as previously described (13, 34). DNA sequencing used this primer pair and also the fungus-specific, universal primer pair ITS1 (5'-TCC GTA GGT GAA CCT GCG G-3') (SEQ ID NO: 62) and ITS4 to amplify a portion of the 18S rDNA region, the entire 5.8S region, the entire ITS1 and ITS2 regions, and a portion of the 28S rDNA region.

A DNA reagent kit (TaKaRa Biomedicals, Shiga, Japan) was used for PCR amplification of genomic DNA. PCR was performed using 2 µl of test sample in a total PCR reaction volume of 100 µl consisting of 10 µl of 10X Ex Taq buffer, 2.5 mM each of dATP, dGTP, dCTP, and dTTP, in 8 µl, 0.2 µM of each primer, and 0.5 U of TaKaRa Ex Taq DNA polymerase.

Thirty cycles of amplification were performed in a Perkin-Elmer 9600 thermal cycler (Emeryville, CA) after initial denaturation of DNA at 95°C for 5 minutes. Each cycle consisted of a denaturation step at 95°C for 30 seconds, an annealing step at 58°C for 30 seconds, and an extension step at 72°C for 1 minute. A final extension at 72°C for 5 minutes followed the last cycle. After amplification, samples were stored at -20°C until used.

Table 1
Synthetic Universal Oligonucleotides Used in PCR and Hybridization Analyses

Primers or Probes	Nucleotide Sequence (5' to 3')	Chemistry and Location
ITS3	GCA TCG ATG AAG AAC GCA GC (SEQ ID NO:63)	5.8S rDNA universal 5' primer
ITS4	TCC TCC GCT TAT TGA TAT GC (SEQ ID NO:64)	28S rDNA universal 3' primer
ITS1	TCC GTA GGT GAA CCT GCG G (SEQ ID NO:62)	18S rDNA universal 5' primer

DNA sequencing

Primary DNA amplifications were conducted as described above. The aqueous phase of the primary PCR reaction was purified using QIAquick Spin Columns (Quiagen, Chatsworth, CA). DNA was eluted from each column with 50 µl of heat-sterilized Tris-EDTA buffer (10 mM Tris, 1 mM EDTA, pH 8.0).

Purified DNA was labeled using a dye terminator cycle sequencing kit (ABI PRISM, Perkin Elmer, Foster City, CA). One mix was made for each of the primers so that sequencing could be performed in both the forward and reverse directions. The reaction volume (20 µl) contained 9.5 µl Terminator Premix, 2 µl (1 ng) DNA template, 1 µl primer (3.2 pmol) and 7.5 µl heat-sterilized distilled H₂O. The mixture was then placed into a pre-heated (96°C) Perkin Elmer 9600 thermal cycler for 25 cycles of 96°C

for 10 seconds, 50°C for 5 seconds, 60°C for 4 minutes. The PCR product was then purified before sequencing using CentriSep spin columns (Princeton Separations, Adelphia, NJ). DNA was then vacuum dried, resuspended in 6 µl of formamide-EDTA (5 µl deionized formamide plus 1 µl 50 mM EDTA, pH 8.0), and denatured for 2 min at 90°C prior to sequencing using an automated capillary DNA sequencer (ABI Systems, Model 373, Bethesda, MD).

The sequencing results were as follows:

Aspergillus flavus 5.8S ribosomal RNA gene, partial sequence, internal transcribed spacer 2, complete sequence, and 28S ribosomal RNA gene, partial sequence.

```
GCTGCCCATC  AAGCACGGC  TTGTGTGTTG  GGTCGTCGTC
CCCTCTCCGG  GGGGGACGGG  CCCCAAAGGC  AGCGGCGGCA
CCGCGTCCGA  TCCTCGAGCG  TATGGGGCTT  TGTCACCCGC
TCTGTAGGCC  CGGCCGGCGC  TTGCCGAACG  CAAATCAATC
TTTTTCCAGG  TTGACCTCGG  ATCAGGTAGG  GATACCCGCT
GAACTTCAA (SEQ ID NO:1)
```

Aspergillus fumigatus 5.8S ribosomal RNA gene, partial sequence, internal transcribed spacer 2, complete sequence, and 28S ribosomal RNA gene, partial sequence.

```
AAACTTTCAA  CAATGGATCT  CTTGGTTCCG  GCATCGATGA
AGAACGCAGC  GAAATGCGAT  AACTAATGTG  AATTGCAGAA
TTCAGTGAAT  CATCGAGTCT  TTGAACGCAC  ATTGCGCCCC
CTGGTATTCC  GGGGGGCATG  CCTGTCCGAG  CGTCATTGCT
GCCCATCAAG  CACGGCTTGT  GTGTTGGGCC  CCCGTCCCCC
TCTCCCGGGG  GACGGGCCCC  AAAGGCAGCG  GCGGCACCGC
GTCCGGTCCT  CGAGCGTATG  GGGCTTGTCA  CCTGCTCTGT
AGGCCCGGCC  GGCGCCAGCC  GACACCCAAC  TTTATTTTTC
TAAGGTTGAC  CTCGGATCAG  GTAGGGATAC  CCGCTGAAC TAAA
(SEQ ID NO:2)
```

Aspergillus niger 5.8S ribosomal RNA gene, partial sequence, internal transcribed spacer 2, complete sequence, and 28S ribosomal RNA gene, partial sequence.

```
AAACTTTCAA  CAATGGATCT  CTTGGTTCCG  GCATCGATGA
AGAACGCAGC  GAAATGCGAT  AACTAATGTG  AATTGCAGAA
```

TTCAGTGAAT CATCGAGTCT TTGAACGCAC ATTGCGCCCC
CTGGTATTCC GGGGGGGCATG CCTGTCCGAG CGTCATTGCT
GCCCTCAAGC ACGGCTTGTG TGTTGGGTCG CCGTCCCCCT
CTCCCGGGGG ACGGGCCCGA AAGGCAGCGG CGGCACCGCG
5 TCCGATCCTC GAGCGTATGG GGCTTTGTCA CCTGCTCTGT
AGGCCCCGGCC GGCGCCTGCC GACGTTATCC AACCATTTTT
TTCCAGGTTG ACCTCGGATC AGGTAGGGAT ACCCGCTGAA CTAA
(SEQ ID NO:3)

10 *Aspergillus terreus* 5.8S ribosomal RNA gene, partial sequence, internal
transcribed spacer 2, complete sequence, and 28S ribosomal RNA gene, partial
sequence.

AAACTTTCAA CAATGGATCT CTTGGTTCCG GCATCGATGA
AGAACGCAGC GAAATGCGAT AACTAATGTG AATTGCAGAA
15 TTCAGTGAAT CATCGAGTCT TTGAACGCAC ATTGCGCCCC
CTGGTATTCC GGGGGGGCAT GCCTGTCCGA GCGTCATTGC
TGCCCTCAAG CCCGGCTTGT GTGTTGGGCC CTCGTCCCCC
GGCTCCCGGG GGACGGGCCC GAAAGGCAGC GGCGGCACCG
CGTCCGGTCC TCGAGCGTAT GGGGCTTCGT CTTCCGCTCC
20 GTAGGCCCGG CCGGCGCCCC CCGAACGCAT TTATTTGCAA
CTTGTTTTTT TTTCCAGGTT GACCTCGGAT CAGGT (SEQ
ID NO:4)

25 *Aspergillus nidulans* 5.8S ribosomal RNA gene, partial sequence, internal
transcribed spacer 2, complete sequence, and 28S ribosomal RNA gene, partial
sequence.

AAACTTTCAA CAATGGATCT CTTGGTTCCG GCATCGATGA
AGAACGCAGC GAACTGCGAT AAGTAATGTG AATTGCAGAA
TTCAGTGAAT CATCGAGTCT TTGAACGCAC ATTGCGCCCC
30 CTGGCATTCC GGGGGGGCATG CCTGTCCGAG CGTCATTGCT
GCCCTCAAGC CCGGCTTGTG TGTTGGGTCG TCGTCCCCC
CCCCGGGGGA CGGGCCCGAA AGGCAGCGGC GGCACCGGTC
CGGTCCTCGA GCGTATGGGG CTTGGTCACC CGCTCGATTA
GGGCCGGCCG GGCGCCAGCC GCGTCTCCA ACCTTATCTT
35 TCTCAGGTTG ACCTCGGATC AGGTAGGGAT ACCCGCTGAA CTAA
(SEQ ID NO:5)

Fusarium solani (strain ATCC62877) internal transcribed spacer 2 and adjacent regions.

5 GAAAATGCGA TAAGTAATGT GAATTGCAGA ATTCAGTGAA
TCATCGAATC TTTGAACGCA CATTGCGCCC GCCAGTATTC
TGGCGGGCAT GCCTGTTCGA GCGTCATTAC AACCCCTCAGG
CCCCCGGGCC TGGCGTTGGG GATCGGCGGA AGCCCCCTGC
GGGCACAACG CCGTCCCCCA AATACAGTGG CGGTCCCGCC
GCAGCTTCCA TTGCGTAGTA GCTAACACCT CGCAACTGGA
10 GAGCGGCGCG GCCACGCCGT AAAACACCCA ACTTCTGAAT
GTTGACCTCG AATCAGGTAG GAATACCCGC TGA ACTTAA (SEQ ID
NO:6)

Fusarium moniliforme (strain ATCC38519) internal transcribed spacer 2 and adjacent regions.

15 AAATGCGATA AGTAATGTGA ATTGCAAAAT TCAGTGAATC
ATCGAATCTT TGAACGCACA TTGCGCCCGC CAGTATTCTG
GCGGGCATGC CTGTTTCGAGC GTCATTTCAA CCCTCAAGCC
CCCGGGTTTG GTGTTGGGGA TCGGCAAGCC CTTGCGGCAA
GCCGGCCCCG AAATCTAGTG GCGGTCTCGC TGCAGCTTCC
20 ATTGCGTAGT AGTAAAACCC TCGCAACTGG TACGCGGCGC
GGCCAAGCCG TTAAACCCCC AACTTCTGAA TGTTGACCTC
GGATCAGGTA GGAATACCCG CTGAACTTAA (SEQ ID NO:7)

Mucor rouxii (strain ATCC24905) internal transcribed spacer 2 and adjacent regions.

25 AAAGTGCAT AACTAGTGTG AATTGCATAT TCAGTGAATC
ATCGAGTCTT TGAACGCAAC TTGCGCTCAT TGGTATTCCA
ATGAGCACGC CTGTTTCAGT ATCAAACAA ACCCTCTATC
CAGCATTTTG TTGAATAGGA ATACTGAGAG TCTCTTGATC
30 TATTCTGATC TCGAACCTCT TGAAATGTAC AAAGGCCTGA
TCTTGTTTAA ATGCCTGAAC TTTTTTTTAA TATAAAGAGA
AGCTCTTGCG GTAAACTGTG CTGGGGCCTC CCAAATAATA
CTCTTTTAA ATTTGATCTG AAATCAGGCG GGATTACCCG
CTGAACTTAA (SEQ ID NO:8)

35 *Mucor racemosus* (strain ATCC22365) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AACTAGTGTG AATTGCATAT TCAGTGAATC
 ATCGAGTCTT TGAACGCAAC TTGCGCTCAT TGGTATTCCA
 ATGAGCACGC CTGTTTCAGT ATCAAAACAA ACCCTCTATC
 CAACTTTTGT TGTATAGGAT TATTGGGGGC CTCTCGATCT
 5 GTATAGATCT TGAAATCCCT GAAATTTACT AAGGCCTGAA
 CTTGTTTAAA TGCCTGAACT TTTTTTTAAT ATAAAGGAAA
 GCTCTTGTA TTTGACTTTGA TGGGGCCTCC CAAATAAATC
 TCTTTTAAAT TTGATCTGAA ATCAGGCGGG ATTACCCGCT
 GAACTTAA (SEQ ID NO:9)

10 *Mucor plumbeus* (strain ATCC4740) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AACTAGTGTG AATTGCATAT TCAGTGAATC
 ATCGAGTCTT TGAACGCAAC TTGCGCTCAT TGGTATTCCA
 15 ATGAGCACGC CTGTTTCAGT ATCAAAACAA ACCCTCTATC
 CAACTTTTGT TGTATAGGAT TATTGGGGGC CTCTCGATCT
 GTATAGATCT TGAAACCCTT GAAATTTACT AAGGCCTGAA
 CTTGTTTAAAT GCCTGAACTT TTTTTTAATA TAAAGGAAAG
 CTCTTGTAAT TGACTTTGAT GGGGCCTCCC AAATAAATCT
 20 TTTTAAATT TGATCTGAAA TCAGGTGGGA TTACCCGCTG
 AACTTAA (SEQ ID NO:10)

Mucor indicus (strain ATCC4857) internal transcribed spacer 2 and adjacent regions.

25 AAAGTGCGAT AACTAGTGTG AATTGCATAT TCAGTGAATC
 ATCGAGTCTT TGAACGCATC TTGCACTCAA TGGTATTCCA
 TTGAGTACGC CTGTTTCAGT ATCAAAAAC AACCCTTATT
 CAAAATTCTT TTTTGAATA GATATGAGTG TAGCAACCTT
 ACAAGTTGAG ACATTTTAAA TAAAGTCAGG CCATATCGTG
 30 GATTGAGTGC CGATACTTTT TTAATTTTGA AAAGGTAAAG
 CATGTTGATG TCCGCTTTTT GGGCCTCCCA AATAACTTTT
 TAAACTTGAT CTGAAATCAG GTGGGATTAC CCGCTGAACT
 TAA (SEQ ID NO:11)

35 *Mucor circinelloides f. circinelloides* (strain ATCC1209B) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AACTAGTGTG AATTGCATAT TCAGTGAATC
 ATCGAGTCTT TGAACGCAAC TTGCGCTCAT TGGTATTCCA
 ATGAGCACGC CTGTTTCAGT ATCAAAACAA ACCCTCTATC
 CAACATTTTT GTTGAATAGG ATGACTGAGA GTCTCTTGAT
 5 CTATTCTGAT CTCGAAGCTC TTGAAATGTA CAAAGGCCTG
 ATCTTGTTTG AATGCCTGAA CTTTTTTTAA ATATAAAGAG
 AAGCTCTTGC GGTAAGCTGT GCTGGGGCCT CCCAAATAAC
 ACATCTTTAA ATTTGATCTG AAATCAGGT GGGACTACCC
 GCTGAACTT AA (SEQ ID NO:12)

10 *Rhizopus oryzae* (strain ATCC34965) internal transcribed spacer 2 and adjacent regions.

AGTGCGATAA CTAGTGTGAA TTGCATATTC AGTGAATCAT
 CGAGTCTTTG AACGCAGCTT GCACTCTATG GTTTTTCTAT
 15 AGAGTACGCC TGCTTCAGTA TCATCACAAA CCCACACATA
 ACATTTGTTT ATGTGGTGAT GGGTCGCATC GCTGTTTTAT
 TACAGTGAGC ACCTAAAATG TGTGTGATTT TCTGTCTGGC
 TTGCTAGGCA GGAATATTAC GCTGGTCTCA GGATCTTTTT
 TTTTGGTTCG CCCAGGAAGT AAAGTACAAG AGTATAATCC
 20 AGTAACTTTC AAACATATGAT CTGAAGTCAG GTGGGATTAC
 CCGCTGAACT TAA (SEQ ID NO:13)

Rhizopus oryzae (strain ATCC11886) internal transcribed spacer 2 and adjacent regions.

25 AGTGCGATAA CTAGTGTGAA TTGCATATTC AGTGAATCAT
 CGAGTCTTTG AACGCAGCTT GCACTCTATG GTTTTTCTAT
 AGAGTACGCC TGCTTCAGTA TCATCACAAA CCCACACATA
 ACATTTGTTT ATGTGGTAAT GGGTCGCATC GCTGTTTTAT
 TACAGTGAGC ACCTAAAATG TGTGTGATTT TCTGTCTGGC
 30 TTGCTAGGCA GGAATATTAC GCTGGTCTCA GGATCTTTTT
 CTTTGGTTCG CCCAGGAAGT AAAGTACAAG AGTATAATCC
 AGCAACTTTC AAACATATGAT CTGAAGTCAG GTGGGATTAC
 CCGCTGAACT TAA (SEQ ID NO:14)

35 *Rhizopus microsporus* (strain ATCC14056) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AACTAGTGTG AATTGCATAT TCGTGAATCA
 TCGAGTCTTT GAACGCAGCT TGC ACTCTAT GGATCTTCTA
 TAGAGTACGC TTGCTTCAGT ATCATAACCA ACCCACACAT
 AAAATTTATT TTATGTGGTG ATGGACAAGC TCGGTAAAT
 5 TTAATTATTA TACCGATTGT CTAAAATACA GCCTCTTTGT
 AATTTTCATT AAATTACGAA CTACCTAGCC ATCGTGCTTT
 TTTGGTCCAA CCAAAAAACA TATAATCTAG GGGTTCTGCT
 AGCCAGCAGA TATTTTAATG ATCTTTAACT ATGATCTGAA
 10 GTCAAGTGGG ACTACCCGCT GAACTTAA (SEQ ID NO:15)

Rhizopus microsporus (strain ATCC12276) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AACTAGTGTG AATTGCATAT TCGTGAATCA
 TCGAGTCTTT GAACGCAGCT TGC ACTCTAT GGATCTTCTA
 15 TAGAGTACGC TTGCTTCAGT ATCATAACCA ACCCACACAT
 AAAATTTATT TTATGTGGTG ATGGACAAGC TCGGTAAAT
 TTAATTATTA TACCGATTGT CTAAAATACA GCCTCTTTGT
 AATTTTCATT AAATTACGAA CTACCTAGCC ATCGTGCTTT
 TTTGGTCCAA CCAAAAAACA TATAATCTAG GGGTTCTGCT
 20 AGCCAGCAAA TATTTTAATG ATCTTTAACC TATGATCTGA
 AGTCAAGTGG GACTACCCGC TGA ACTTAA (SEQ ID NO:16)

Rhizopus circinans (strain ATCC34106) internal transcribed spacer 2 and adjacent regions.

AAATTGCGAT AACTAGTGTG AATTGCATTT TCAGTGAATC
 ATCGAGTCTT TGAACGCAT CTTGCGCTCT TGGGATTCTT
 CCCTAGAGCA CACTTGCTTC AGTATCATAA CAAAACCCTC
 ACCTAATATT TTTTTTTTTT AAAAAAAAAA TATTAGAGTG
 GTATTGGGGT CTCTTTGGTA ATTCTTTGTA ATTATAAAAG
 30 TACCCTTAAA TGTCATAAAC AGGTTAGCTT TAGCTTGCCT
 TTAAAGATCT TCTTAGGGTA TCATTACTTT TCGTAAATCT
 TTAATAGGCC TGTCACATAA TTCTACCCTT AAATTTCTTA
 AACCTTGATC TGAAGTCAAG TGGGAGTACC CGCTGAACTT AA
 (SEQ ID NO:17)

Rhizopus circinans (strain ATCC34101) internal transcribed spacer 2 and adjacent regions.

```

5  AAATTGCGAT  AACTAGTGTG  AATTGCATTT  TCAGTGAATC
   ATCGAGTCTT  TGAACGCATC  TTGCGCTCTT  GGGATTCTTC
   CCTAGAGCAC  ACTTGCTTCA  GTATCATAAC  AAAACCCTCA
   CCTAATATTT  TTTTAAATA  AAAAAAATA  TTAGAGTGGT
   ATTGGGGTCT  CTTTGGTAAT  TCTTTGTAAT  TATAAAAGTA
   CCCTTAAATG  TCATAAACAG  GTTAGCTTTA  GCTTGCCTTT
10  AAAGATCTTC  TTAGGGTATC  ATTACTTTTC  GTAAATCTTT
   AATAGGCCTG  TCACATAATT  CTACCCTTAA  ATTTCTTAAA
   CCTTGATCTG  AAGTCAAGTG  GGAGTACCCG  CTGAACTTAA (SEQ
   ID NO:18)

```

15 *Rhizopus stolonifer* (strains ATCC14037 and 6227A) internal transcribed spacer 2 and adjacent regions.

```

   AAAGTGCGAT  AACTAGTGTG  AATTGCATAT  TCAGTGAATC
   ATCGAGTCTT  TGAACGCAAC  TTGCACTCTA  TGGTTTTCCG
   TAAAGTACGC  TTGCTTCAGT  ATCATAAAGA  CCCCATCCTG
   ATTATTATTT  TTTTATTAAT  ATAATTAATT  TTGGAGATAA
20  TAAAAATGAG  GCTCTTTCTT  TTCTTTTTTT  TTTTTTAAAT
   AAAAAGGGGG  GGAAAGGGTC  TTTTAAATG  GGCAAATTCT
   GGGTTTTTTA  CTAAACCTGA  ACTCCCCCA  AAAATTCAAA
   AAAAAAAAAA  TGGGTTTTAC  CAAATTTTTT  TTTTTTTTCT
   CCTTTTTGTG  TAGTTAATAC  TCTATTAAAT  TTATTTACTT
25  GGTATTATAA  CGATTATGCA  AGAAGGGAGA  GAACAAAGAA
   TAATGAAAGA  GAGTTTTTAA  ATAAATTCTT  TTTTCATTTT
   TTCAATCAAT  GATCTGAAGT  CAAGTGGGAT  TACCCGCTGA
   ACTTAA (SEQ ID NO:19)

```

30 *Rhizomucor pusillus* (strain ATCC36606) internal transcribed spacer 2 and adjacent regions.

```

   AAATTGCGAA  AAGTAATGCG  ATCTGCAGCC  TTTGCGAATC
   ATCGAATTCT  CGAACGCACC  TTGCACCCTT  TGGTTCATCC
   ATTGGGTACG  TCTAGTTCAG  TATCTTTATT  AACCCTAAA
35  GGTTTATTTT  TTGATAAATC  TTTGGATTG  CGGTGCTGAT
   GGATTTTCAT  CCGTTCAAGC  TACCCGAACA  ATTTGTATGT
   TGTTGACCCT  TGATATTTC  TTGAGGGCTT  GCATTGGTAT

```

CTAATTTTTT ACCAGTGTGC TTCGAGATGA TCAAGTATAA
 AGGTCAATCA ACCACAAATA AATTTCAACT ATGGATCTGA
 ACTTAGATGG GATTACCCGC TGAACCTAA (SEQ ID NO:20)

5 *Absidia corymbifera* (strain ATCC46774) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AATTATTGCG ACTTGCATTC ATAGCGAATC
 ATCGAGTTCT CGAACGCATC TTGCGCCTAG TAGTCAATCT
 ACTAGGCACA GTTGTTTCAG TATCTGCAAC TACCAATCAG
 10 TTCAACTTGG TTCTTTGAAC CTAAGCGAGC TGGAAATGGG
 CTTGTGTTGA TGGCATTTCAG TTGCTGTCAT GGCCTTAAAT
 ACATTTAGTC CTAGGCAATT GGCTTTAGTC ATTTGCCGGA
 TGTAGACTCT AGAGTGCCTG AGGAGCAACG ACTTGGTAG
 TGAGTTCATA ATTCCAAGTC AATCAGTCTC TTCTTGAAC
 15 AGGTCTTAAT CTTTATGGAC TAGTGAGAGG ATCTAACTTG
 GGTCTTCTCT TAAAACAAAC TCACATCTAG ATCTGAAATC
 AACTGAGATC ACCCGCTGAA CTAA (SEQ ID NO:21)

20 *Absidia corymbifera* (strain ATCC46773) internal transcribed spacer 2 and adjacent regions.

AAAGTGCGAT AATTATTGCG ACTTGCATTC ATAGTGAATC
 ATCGAGTTCT TGAACGCATC TTGCGCCTAG TAGTCAATCT
 ACTAGGCACA GTTGTTTCAG TATCTGCATC CACCAATCAA
 CTTAACCTTT TGTGTTGAGT TGGAACCTGGG CTTCTAGTTG
 25 ATGGCATTTA GTTGCTGTCA TGGCCTTAAA TCAATGTCCT
 AGGTGTTAGA ACATCTAACA CCGGATGGAA ACTTTAGAGC
 GCTTTAAGAG CAGCTTGGTT AGTGAGTTCA ATAATTCCAA
 GCATTAAGTC TTTTAATGAA CTAGCTTTTC TATCTATGGG
 AACTACTTG GAGAAATCCA AGTAACCTTT AAACCTCCAT
 30 TTAGATCTGA AATCAACTGA GACCACCCGC TGAACCTAA (SEQ ID NO:22)

Cunninghamella elegans (strain ATCC42113) internal transcribed spacer 2 and adjacent regions.

35 AAATCGCGAT ATGTAATGTG ACTGCCTATA GTGAATCATC
 AAATCTTTGA AACGCATCTT GCACCTTATG GTATTCCATA
 AGGTACGTCT GTTTCAGTAC CACTAATAAA TCTCTCTCTA

TCCTTGATGA TAGAAAAAAA AAAAATAATT TTTACTGGGC
CCGGGGAATC CTTTTTTTTT TTTAATAAAA AGGACCAATT
TTGGCCCAAA AAAAAGGGTT GAACTTTTTT TACCAGATCT
TGCATCTAGT AAAAACCTAG TCGGCTTTAA TAGATTTTAA
5 TTTTCTATTA AGTTTATAGC CATTCTTATA TTTTTTAAAA
TCTTGGCCTG AAATCAGATG GGATACCCGC TGAACCTAA (SEQ ID
NO:23)

Pseudallescheria boydii (strain ATCC44328) internal transcribed spacer 2 and
adjacent regions (teleomorph of *Scedosporium apiospermum*).

10 AAATGCGATA AGTAATGTAA ATTGCAAAAT TCAGTGAATC
ATCGAATCTT TGAAACGCAC ATTGCGCCCG GCAGTAATCT
GCCGGGCATG CCTGTCCGAG CGTCATTTC ACCCTCGAAC
CTCCGTTTC CTTAGGGAAG CCTAGGGTCG GTGTTGGGGC
15 GCTACGGCAA GTCCTCGCAA CCCCCGTAGG CCCTGAAATA
CAGTGGCGGT CCCGCCGCGG TTGCCTTCTG CGTAGTAAGT
CTCTTTTGCA AGCTCGCATT GGGTCCCGGC GGAGGCCTGC
CGTCAAACCA CCTAACAACCT CCAGATGGTT TGACCTCGGA
TCAGGTAGGG TTACCCGCTG AACTTAA (SEQ ID NO:24)

20 *Pseudallescheria boydii* (strain ATCC36282) internal transcribed spacer 2 and
adjacent regions (teleomorph of *Scedosporium apiospermum*).

GAAATGCGAT AAGTAATGTG AATTGCAGAA TTCAGTGAAT
CATCGAATCT TTGAAACGCA CATTGCGCCC GGCAGTAATC
25 TGCCGGGCAT GCCTGTCCGA GCGTCATTTC AACCCTCGAA
CCTCCGTTTC CTCAGGGAAG CTCAGGGTCG GTGTTGGGGC
GCTACGGCAA GTCTTCGCAA CCCTCCGTAG GCCCTGAAAT
ACAGTGGCGG TCCCGCCGCG GTTGCCTTCT GCGTAGAAGT
CTCTTTTGCA AGCTCGCATT GGGTCCCGGC GGAGGCCTGC
30 CGTCAAACCA CCTATAACTC CAAATGGTTT GACCTCGGAT
CAGGTAGGGT TACCCGCTGA ACTTAA (SEQ ID NO:25)

Scedosporium apiospermum (strain ATCC64215) internal transcribed spacer 2
and adjacent regions.

35 GAAATGCGAT AAGTAATGTG AATTGCAGAA TTCAGTGAATC
ATCGAATCTT TGAACGCACA TTGCGCCCGG CAGTAATCTG
CCGGGCATGC CTGTCCGAGC GTCATTTC AA CCCTCGAACC

5 TCCGTTTCCT CAGGGAAGCT CAGGGTCGGT GTTGGGGGCGC
TACGGCGAGT CTTCGCGACC CTCCGTAGGC CCTGAAATAC
AGTGGCGGTC CCGCCGCGGT TGCCTTCTGC GTAGTAAGTC
TCTTTTGCAA GCTCGCATTG GGTCCCGGCG GAGGCCTGCC
GTCAAACCAC CTATAACTCC AGATGGTTTG ACCTCGGATC
AGGTAGGTAC CCGCTGAACT TAA (SEQ ID NO:26)

Scedosporium apiospermum (strain ATCC46173) internal transcribed spacer 2 and adjacent regions.

10 AAATGCGATA AGTAATGTGA ATTGCAGAAT TCAGTGAATC
ATCGAATCTT TGAACGCACA TTGCGCCCGG CAGTAATCTG
CCGGGCATGC CTGTCCGAGC GTCATTTCOA CCCTCGAACC
TCCGTTTCCT CAGGGAAGCT CAGGGTCGGT GTTGGGGGCGC
TACGGCGAGT CTTCGCGACC CTCCGTAGGC CCTGAAATAC
15 AGTGGCGGTC CCGCCGCGGT TGCCTTCTGC GTAGTAAGTC
TCTTTTGCAA GCTCGCATTG GGTCCCGGCG GAGGCCTGCC
GTCAAACCAC CTATAACTCC AGATGGTTTG ACCTCGGATC
AGGTAGGTAC CCGCTGAACT TAA (SEQ ID NO:27)

20 *Penicillium notatum* (strain ATCC10108) internal transcribed spacer 2 and adjacent regions.

AAATGCGATA CGTAATGTGA ATTGCAAATT CAGTGAATCA
TCGAGTCTT TGAACGCACA TTGCGCCCCC TGGTATTCCG
GGGGGCATGC CTGTCCGAGC GTCATTGCTG CCCTCAAGCA
25 CGGCTTGTGT GTTGGGGCCCC GTCCTCCGAT CCCGGGGGAC
GGGCCCCGAAA GGCAGCGGCG GCACCGCGTC CGGTCCTCGA
GCGTATGGGG CTTTGTACAC CGCTCTGTAG GCCCGGCCGG
CGCTTGCCGA TCAACCCAAA TTTTATCCA GGTGACCTC
GGATCAGGTA GGGATACCCG CTGAACTTAA (SEQ ID NO:28)

30 *Sporothrix schenckii* (strain ATCC14284) internal transcribed spacer 2 and adjacent regions.

GAAATGCGAT ACTAATGTGA ATTGCAGAAT TCAGCGAACC
ATCGAATCTT TGAACGCACA TTGCGCCCGC CAGCATTCTG
35 GCGGGCATGC CTGTCCGAGC GTCATTTCCT CCCTCACGCG
CCCCGTTGCG CGCTGGTGTT GGGGCGCCCT CCGCCTGGCG
GGGGGCCCCC GAAAGCGAGT GGCGGGCCCT GTGGAAGGCT

CCGAGCGCAG TACCGAACGC ATGTTCTCCC CTCGCTCCGG
AGGCCCCCCA GGCGCCCTGC CGGTGAAAAC GCGCATGACG
CGCAGCTCTT TTTACAAGGT TGACCTCGGA TCAGGTGAGG
ATACCCGCTG ACTTAA (SEQ ID NO:29)

Contamination precautions

Precautions were taken to avoid possible contamination of PCR samples by following the guidelines of Fujita and Kwok (13, 22). All buffers and distilled water used for PCR assays were autoclaved and fresh PCR reagents were aliquoted prior to use. Physical separation of laboratory areas used to prepare PCR assays and to analyze PCR products, and the use of aerosol-resistant pipette tips, reduced possible cross-contamination of samples by aerosols. Appropriate negative controls were included in each test run, including controls omitting either the primer or the DNA template during PCR assays.

Agarose gel electrophoresis

Gel electrophoresis was conducted in TBE buffer (0.1 M Tris, 0.09 M boric acid, 1 mM EDTA, pH 8.4) at 80 V for 1 to 2 hours using gels composed of 1% (w/vol) agarose (International Technologies, New Haven, CT) and 1% (w/vol) NuSieve agar (FMC Bioproducts, Rockland, ME). Gels were stained with 0.5 µg of ethidium bromide (EtBr) per ml of distilled H₂O for 10 minutes followed by three serial washes for 10 minutes each with distilled H₂O.

Microtitration plate enzyme immunoassay for the detection of PCR products

Amplicons were detected using species-specific and genus probes labeled with digoxigenin and an all-filamentous fungal probe labeled with biotin in a streptavidin-coated microtiter plate format (13, 34). Ten µl of PCR product was added to each 1.5 ml Eppendorf tube. Single-stranded DNA was then prepared by heating the tubes at 95°C for 5 minutes and cooling immediately on ice. Two-tenths of a ml of hybridization solution [4x SSC (saline sodium citrate buffer, 0.6 M NaCl, 0.06 M trisodium citrate, pH 7.0) containing 20 mM Hepes, 2 mM EDTA, and 0.15% (vol/vol) Tween 20] supplemented with 50 ng/ml each of the all-*Aspergillus* biotinylated probe and a species-specific digoxigenin-labeled probe was added to each

5 tube containing denatured PCR product. Tubes were mixed by inversion and placed in a water bath at 37°C to allow probes to anneal to PCR product DNA. After 1 hour, 100 µl of each sample was added to duplicate wells of a commercially prepared streptavidin-coated microtitration plate (Boehringer
10 Mannheim, Indianapolis, IN). The plate was incubated at ambient temperature for 1 hour with shaking, using a microtitration plate shaker (manufactured for Dynatech by CLTI, Middletown, NY). Plates were washed 6 times with 0.01 M potassium phosphate buffered saline, pH 7.2, containing 0.05% Tween 20 (PBST). Each well then received 100 µl of
15 horseradish peroxidase-conjugated, anti-digoxigenin Fab fragment (Boehringer Mannheim) diluted 1:1000 in hybridization buffer. After incubation at ambient temperature for 30 minutes with shaking, the plate was washed 6 times with PBST. One hundred µl of a mixture of one volume of 3, 3', 5, 5'-tetramethyl benzidine peroxidase substrate (Kirkegaard and
20 Perry Laboratories, Inc., Gaithersburg, MD) and one volume of peroxidase solution (Kirkegaard and Perry Laboratories) was added to each well and the plate was placed at ambient temperature for 10 minutes for color development. The $A_{650\text{nm}}$ of each well was determined with a microtitration plate reader (UV Max, Molecular Devices, Inc., Menlo Park, CA). The absorbance value for the reagent blank, where DNA was absent but replaced with distilled H₂O, was subtracted from each test sample.

Statistical analysis

25 The Student's t test was used to determine differences between sample means. Means are expressed as the mean plus or minus the standard error from the mean. Differences were considered significant when $P < 0.05$.

30 The following probes were used to detect and distinguish each species.

Table 2
Probe Sequences

PROBES	5' to 3' OLIGONUCLEOTIDE SEQUENCE	
Generic Biotin Probe	5' end-labeled biotinylated probe 5.8S region of rDNA	
B-58	GAA TCA TCG A(AG)T CTT TGA ACG	SEQ ID NO 61
Digoxigenin-probe	5' end-labeled digoxigenin probe ITS2 region of rDNA	
Aspergillus species		
A. flavus 22	GCA AAT CAA TCT TTT TCC	SEQ ID NO 30
A. flavus 23	GAA CGC AAA TCA ATC TTT	SEQ ID NO 31
A. fumigatus	CCG ACA CCC ATC TTT ATT	SEQ ID NO 32
A. niger	GAC GTT ATC CAA CCA TTT	SEQ ID NO 33
A. nidulans	GGC GTC TCC AAC CTT ATC	SEQ ID NO 35
A. terreus	GCA TTT ATT TGC AAC TTG	SEQ ID NO 34
Fusarium species		
F. moniliforme	TCT AGT GAC GGT CTC GCT	SEQ ID NO 49
F. oxysporum	CGT TAA TTC GCG TTC CTC	SEQ ID NO 50
F. solani	CTA ACA CCT CGC AAC TGG AGA	SEQ ID NO 51
Mucor species		
M. circinelloides	AAC ATT TTT GTG AAT AGG ATG	SEQ ID NO 39
M. indicus	CGT GGA TTG AGT GCC GAT	SEQ ID NO 38
M. plumbeus	GAA ACC CTT GAA ATT	SEQ ID NO 37
M. rouxii	GAA TAG GAA TAC TGA GAG	SEQ ID NO 36
M. racemosus	GAA ATC CCT GAA ATT	SEQ ID NO 40
Penicillium species		

Penicillium marneffei 1	GGG TTG GTC ACC ACC ATA	SEQ ID NO 47
Penicillium marneffei 2	TGG TCA CCA CCA TAT TTA	SEQ ID NO 48
Penicillium notatum	GAT CAA CCC AAA TTT TTA	SEQ ID NO 46
Rhizopus species		
R. circinans	CTT AGG GTA TCA TTA CTT	SEQ ID NO 42
R. microsporus	CAT ATA ATC TAG GGG TTC	SEQ ID NO 57
R. oryzae	GAG TAT AAT CCA G(CT)A ACT	SEQ ID NO 41
R. stolonifer	CTT GGT ATT ATA ACG ATT	SEQ ID NO 44
Rhizomucor pusillus	TCC TTG AGG GCT TGC ATT	SEQ ID NO 43
Other Genera		
Absidia corymbifera	GTT GCT GTC ATG GCC TTA	SEQ ID NO 55
Cunninghamella elegans 4	TAG TCG GCT TTA ATA GAT	SEQ ID NO 52
Cunninghamella elegans 5	TAT TAA GTT TAT AGC CAT	SEQ ID NO 53
Cunninghamella elegans 6	TAA GTT TAT AGC CAT TCT	SEQ ID NO 54
Pseudallescheria boydii	AAG TCT CTT TTG CAA GCT	SEQ ID NO 45
Sporothrix schoenckii	GAC GCG CAG CTC TTT TTA	SEQ ID NO 56
Genus Probes		
G-ASPERGILLUS	CCT CGA GCG TAT GGG GCT	SEQ ID NO 58
G-FUSARIUM	CCC AAC TTC TGA ATG TTG	SEQ ID NO 59
G-MUCOR	(AC)TG GGG CCT CCC AAA TAA	SEQ ID NO 60

Species-specific probes to the ITS2 region of rDNA for *Aspergillus fumigatus* (SEQ ID NO:32), *A. flavus* (SEQ ID NO:31), *A. niger* (SEQ ID NO:33), *A. terreus* (SEQ ID NO:34), and *A. nidulans* (SEQ ID NO:35) correctly identified each of the respective species ($P < 0.001$), and gave no false-positive reactions with *Rhizopus*, *Mucor*, *Fusarium*, *Penicillium*, or *Candida* species. The *A. flavus* probe also recognized *A. oryzae*, which belongs to the *A. flavus* group. Identification time was reduced from a mean of 5 days by conventional methods to 8 hours.

Table 3
***Aspergillus* Probes**

Fungus	<i>A. fumigatus</i>	<i>A. nidulans</i>	<i>A. niger</i>	<i>A. terreus</i>	<i>A. flavus</i>
<i>A. fumigatus</i> (n=6)	2.197 ± 0.187	0.002	0.000	0.001	0.001
<i>A. nidulans</i> (n=3)	0.001	1.315 ± 0.464	0.002	0.000	0.001
<i>A. niger</i> (n=5)	0.000	0.000	1.242 ± 0.471	0.001	0.003
<i>A. terreus</i> (n=4)	0.001	0.000	0.001	1.603 ± 0.378	0.001
<i>A. flavus</i> (n=6)	0.001	0.001	0.000	0.001	2.043 ± 0.390
<i>A. oryzae</i> (n=2)	0.001	0.002	0.001	0.001	2.445 ± 0.106
<i>A. parasitica</i> (n=1)	0.001	0.002	0.002	0.002	0.051
<i>A. clavus</i> (n=1)	0.005	0.005	0.006	0.005	0.003
<i>C. albicans</i> (n=1)	0.002	0.001	0.002	0.000	0.000
<i>C. parasitosis</i> (n=1)	0.001	0.002	0.002	0.002	0.001
<i>C. glabrata</i> (n=1)	0.001	0.003	0.001	0.001	0.005

C.krusei (n=1)	0.002	0.002	0.002	0.001	0.001
C.tropicalis (n=1)	0.002	0.002	0.001	0.000	0.001
F.moniliforme (n=1)	0.003	0.003	0.001	0.001	0.001
F.solani (n=1)	0.006	0.002	0.001	0.000	0.001
R.oryzae (n=1)	0.001	0.001	0.001	0.001	0.001
M.racemosus (n=1)	0.001	0.002	0.005	0.002	0.000
P.notatum (n=1)	0.001	0.002	0.002	0.002	0.000
Avg±SD negative controls	0.001 ± 0.002	0.001 ± 0.001	0.000 ± 0.002	0.000 ± 0.002	0.002 ± 0.010

Species-specific probes to the ITS2 region of rDNA for *Fusarium oxysporum*, *F. solani*, and *F. moniliforme*, correctly identified each of the respective species ($P < 0.001$), and gave no false-positive reactions with *Blastomyces*, *Apophysomyces*, *Candida*, *Aspergillus*, *Mucor*, *Penicillium*, *Rhizopus*, *Rhizomucor*, *Absidia*, *Cunninghamella*, *Pseudallescheria*, *Sporothrix*, or *Neosartorya*. Empty boxes in Table 4 represent zero probe reactivity.

Table 4
***Fusarium* Probes**

Fungus	F. oxysporum	F. solani	F. moniliforme	Generic Fusarium
F. oxysporum (n=3)	1.40 ± 0.13			1.76 ± 0.27
F. solani (n=5)		1.57 ± 0.07		1.35 ± 0.28

F. moniliforme (n=2)			1.40 ± 0.01	1.34 ± 0.91
Negative control				
Fungus	F. oxysporum	F. solani	F. moniliforme	Generic Fusarium
A.fumigatus				
A.flavus				
A.niger				
A.nidulans				
A.terreus				
A.parasiticus				
A.clavatus				
P.marneffei		0.01	0.01	
P.notatum	0.01	0.01	0.01	
Rhizopus oryzae		0.03	0.01	
Rhizopus microsporus		0.01	0.01	
Rhizopus circinans		0.01	0.01	
Rhizopus stolonifer		0.01	0.01	
Rhizomucor pusillus		0.03	0.02	
M. racemosus				
M. circinelloides				
M. rouxii				
M. plumbeus				
M. indicus				
Absidia corymbifera		0.01	0.01	
Cunninghamella elegans		0.01	0.02	
P. boydii			0.02	
Sporothrix schenckii		0.01	0.01	
C.albicans				
C.tropicalis				
C.krusei				
C.parasilosis				
C.glabrata				

Neosartorya fischeri		0.01		
Blastomyces dermatitidis				
Apophysomyces elegans				
Average of negative controls	0.001 ± 0.002	0.005 ± 0.01	0.004 ± 0.006	

Species-specific probes to various other zygomycetes are presented in Table 5, showing correct identification of each species and no false positives. The exceptions are that the *M. circinelloides* probe hybridized with the *M. rouxii* DNA and the *M. plumbeus* probe hybridized with the *M. racemosus* DNA. However, the *M. rouxii* probe did not hybridize with *M. circinelloides* DNA, nor did the *M. racemosus* probe hybridize with *M. plumbeus* DNA. Therefore, by a process of elimination, each species can be correctly identified. Empty boxes in Table 5 represent zero probe reactivity.

	D- probes	RMIC	RCIR	RSTOL	RPUS	MACE	MCIR	MRX	MPLUM	MIND	ABS	CUN
FUNGUS	RORY 1.50 ± 0.48				0.01							
R. oryzae (n=5)		0.96 ± 0.61										
R. microsporus (n=5)												
R.circinans (n=3)			1.56 ± 0.19									
R. stolonifer (n=5)				2.53 ± 0.07			0.01					
Rhizomucor pusillus (n=2)					1.10 ± 0.68							
M. racemosus (n=6)				0.01		2.02 ± 0.34			0.29 ± 0.52			
M.circineloides (n=3)							1.63 ± 0.37	0.01	0.02			
M. rouxii (n=1)							1.77	0.76				
M.plumbeus (n=2)									2.14 ± 0.25			
M. indicus (n=1)		0.01								1.70 ± 0.04		
Absidia corymbifera (n=2)					0.01				0.01		1.61 ± 0.08	
Cunninghamella elegans (n=2)		0.01										2.26 ± 0.03

Table 5 Continued

Negative control	D- probes RORY	RMIC	RCIR	RSTOL	RPUS	MRACE	MCIR	MRX	MPLUM	MIND	ABS	CUN
FUNGUS												
<i>A.fumigatus</i>									0.01	0.02		
<i>A.flavus</i>					0.01					0.05		
<i>A.niger</i>								0.01				
<i>A.nidulans</i>									0.01	0.01		
<i>A.terreus</i>	0.01											
<i>A.parasiticus</i>					0.01					0.03		
<i>A.clavatus</i>										0.02		
<i>P.marneffei</i>			0.01									
<i>P.notatum</i>								0.01		0.03		
<i>F.oxysporum</i>												
<i>F.solani</i>												
<i>F.moniliforme</i>	0.01			0.01				0.01		0.01		
<i>P.boydii</i>	0.02											
<i>Sporothrix schenckii</i>												
<i>C.albicans</i>												
<i>C.tropicalis</i>												
<i>C.krusei</i>												
<i>C.parasilosis</i>												
<i>C.glabrata</i>												
<i>Neosartorya fischeri</i>			0.01									
<i>Blastomyces dermatitidis</i>												
<i>Apophysomyces elegans</i>												
Average	0.001 ± .004	0.001 ± 0.02	0.000 ± 0.002	0.000 ±0.003	0.001 ±0.003	0.001 ± 0.002	0.001 ±0.002	0.001 ±0.003	0.003 ±0.005	0.005 ± 0.01	0.001 ±0.001	

Species-specific probes to various other fungi are presented in Table 6, showing correct identification of each species and no false positives. Empty boxes in Table 6 represent zero probe reactivity.

5

Table 6
Pseudallescheria and *Sporothrix* Probes

Fungus	<i>P. boydii</i>	<i>P.marneffei</i>	<i>P.notatum</i>	<i>Sporothrix schenckii</i>
<i>P. boydii</i> (n=4)	1.65 ± 0.48			
<i>P.marneffei</i> (n=3)	0.01	1.24 ± 0.12		
<i>P.notatum</i> (n=3)			1.93 ± 0.25	
<i>Sporothrix schenckii</i> (n=3)	0.01			1.94 ± 0.25
Negative control				
Fungus	<i>P. boydii</i>	<i>P.marneffei</i>	<i>P.notatum</i>	<i>Sporothrix schenckii</i>
<i>A.fumigatus</i>	0.01			
<i>A.flavus</i>				
<i>A.niger</i>				
<i>A.nidulans</i>				
<i>A.terreus</i>				
<i>A.parasiticus</i>				
<i>A.clavatus</i>			0.11	
<i>F.oxysporum</i>		0.10		
<i>F. solani</i>		0.14		
<i>F. moniliforme</i>		0.08		
<i>R. oryzae</i>	0.01			
<i>R. microsporus</i>	0.01			
<i>R. circinans</i>	0.01			

R. stolonifer	0.01			
Rhizomucor pusilus				
M. racemosus		0.04		
M. circinelloides	0.01	0.09		
M. rouxii	0.01			
M. plumbeus		0.05		
M. indicus				
Absidia corymbifera	0.01			
Cunninghamella bertholletiae	0.01			
C. albicans				
C. tropicalis		0.02		
C. krusei				
C. parasitosis				
C. glabrata				
Neosartorya pseudofischeri		0.03		
Blastomyces dermatitidis	0.01			
Apophysomyces elegans	0.01			
Average Negative Controls	0.004 ± 0.002	0.013 ± 0.03	0.002 ± 0.019	0.001 ± 0.002

All of the references mentioned in this Specification are hereby incorporated by reference in their entirety.

References

1. **Ampel, N.M.** 1996. Emerging disease issues and fungal pathogens associated with HIV infection. **Emerg. Infect. Dis.** 2:109-116.
2. **Andriole, V.T.** 1996. *Aspergillus* infections: problems in diagnosis and treatment. **Infect. Agents and Dis.** 5:47-54.
3. **Andriole, V.T.** 1993. Infections with *Aspergillus* species. **Clin. Infect. Dis.** 17 Suppl 2:S481-S486.
4. **Bir, N., A. Paliwal, K. Muralidhar, P. Reddy, and P.U. Sarma.** 1995. A rapid method for the isolation of genomic DNA from *Aspergillus fumigatus*. **Prep. Biochem.** 25:171-181.

5. Blum, U., M. Windfuhr, C. Buitrago-Tellez, G. Sigmund, E.W. Herbst, and M. Langer. 1994. Invasive pulmonary aspergillosis. MRI, CT, and plain radiographic findings and their contribution for early diagnosis. *Chest* 106:1156-1161.
6. Caillot, D., O. Casasnovas, A. Bernard, J.F. Couaillier, C. Durand, B. Cuisenier, E. Solary, F. Piard, T. Petrella, A. Bonnin, G. Couaillault, M. Dumas, and H. Guy, 1997. Improved management of invasive pulmonary aspergillosis in neutropenic patients using early thoracic computed tomographic scan and surgery. *J. Clin. Oncol.* 15:139-147.
7. Denning, D.W. Therapeutic outcome in invasive aspergillosis. *Clin. Infect. Dis.* 23:608-615.
8. Denning, D.W. Diagnosis and management of invasive aspergillosis. *Curr. Clin. Topics Inf. Dis.* 16:277-299.
9. de Repentigny, L., L. Kaufman, G. T. Cole, D. Kruse, J. P. Latge, and R. C. Matthews. 1994. Immunodiagnosis of invasive fungal infections. *J. Med. Vet. Mycol.* 32 Suppl 1239-252.
10. Dupont, B., D. W. Denning, D. Marriott, A. Sugar, M. A. Viviani, and T. Sirisanthana. 1994. Mycoses in AIDS patients. *J. Med. Vet. Mycol.* 32 Suppl 1:221-239.
11. Fisher, B. D., D. Armstrong, B. Yu, and J. W. M. Gold. 1981. Invasive aspergillosis: progress in early diagnosis and treatment. *Am. J. Med.* 71:571-577.
12. Fridkin, S. K. and W. R. Jarvis. 1996. Epidemiology of nosocomial fungal infections. *Clin. Microbiol. Rev.* 9:499-511.
13. Fujita, S-I., B.A. Lasker, T. J. Lott, E. Reiss, and C. J. Morrison. 1995. Micro titration plate enzyme immunoassay to detect PCR-amplified DNA from *Candida* species in blood. *J. Clin. Microbiol.* 33:962-967.
14. Gordon, M. A., E. W. Lapa, and J. Kane. 1977. Modified indirect fluorescent antibody test for aspergillosis. *J. Clin. Microbiol.* 6:161-165.

15. **Holmes, A. R., R. D. Cannon, M. G. Shepard, and H. F. Jenkinson.** 1994. Detection of *Candida albicans* and other yeast in blood by PCR. *J. Clin. Microbiol.* **32**:228-231.
- 5 16. **Hung, C. C., S. C. Chang, P. C. Yang, W. C. Hseigh.** 1994. Invasive pulmonary pseudallescheriasis with direct invasion of the thoracic spine in an immunocompromised patient. *Eur. J. Clin. Microbiol. Infect. Dis.* **13**:749-751.
- 10 17. **Kappe, R., and H. P. Seeliger.** 1993. Serodiagnosis of deep-seated fungal infections. *Curr. Topics Med. Mycol.* **5**:247-280.
18. **Kappe, R., A. Schulze-Berge, H. G. Sonntag.** 1996. Evaluation of eight antibody tests and one antigen test for the diagnosis of invasive aspergillosis. *Mycoses* **39**:13-23.
- 15 19. **Kaufman and Reiss,** Manual of Clinical Microbiology.
- 20 20. **Kremery, V., Jr., E. Kunova, Z. Jesenska, J. Trupl, S. Spanik, J. Mardiak, M. Studena, and E. Kukuckova.** 1996. Invasive mold infections in cancer patients: 5 years' experience with *Aspergillus*, *Mucor*, *Fusarium* and *Acremonium* infections. *Supportive Care in Cancer* **4**:39-45.
21. **Khoo, S. H., and D. W. Denning.** 1994. Invasive aspergillosis in patients with AIDS. *Clin. Infect. Dis* **19** Suppl 1: S41-S48.
- 25 22. **Kwok, S., and R. Higuichi.** 1989. Avoiding false positives with PCR. *Nature (London)* **339**:237-238.
23. **Larone, D. H.** Medically Important Fungi: A Guide to Identification. 3rd ed. ASM Press, Washington, D. C. 1995.
- 30 24. **Leenders, A., A. van Belkum, S. Janssen, S. de Marie, J. Kluytmans, J. Wielenga, B. Lowenberg, and H. Verbrugh.** Molecular epidemiology of apparent outbreak of invasive aspergillosis in a hematology ward. *J. Clin. Microbiol.* **34**:345-351.
- 35 25. **Makimura, K., S. Y. Murayama, H. Yamaguchi.** 1994. Specific detection of *Aspergillus* and *Penicillium* species from respiratory specimens by polymerase chain reaction (PCR). *Jap. J. Med. Sci. Biol.* **47**:141-156.

26. **Maniatis, T., E. F. Fritsch, and J. Sambrook.** 1982. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
27. **Martino, P., and C. Girmenia.** 1993. Diagnosis and treatment of invasive fungal infections in cancer patients. Supportive Care in Cancer. 1:240-244.
28. **Melchers, W. J., P. E. Verweij, P. van den Hurk, A. van Belkum, B. E. De Pauw, J. A. Hoogkamp-Korstanje, and J. F. Meis.** 1994. General primer-mediated PCR for detection of *Aspergillus* species. J. Clin. Microbiol. 32:1710-1717.
29. **Miller, W. T. J., G. J. Sals, I. Frank, W. B. Geftter, M. Aronchick, W. T. Miller.** 1994. Pulmonary aspergillosis patients with AIDS. Clinical and radiographic correlations. Chest 105:37-44.
30. **Miyakawa, Y., T. Mabuchi, and Y. Fukazawa.** 1993. New method for detection of *Candida albicans* in human blood by polymerase chain reaction. J. Clin. Microbiol. 31:3344-3347.
31. **Montone, K. T., and L. A. Litzky.** 1995. Rapid method for detection of *Aspergillus* 5S ribosomal RNA using a genus-specific oligonucleotide probe. J. Clin. Microbiol. 103:48-51.
32. **Rogers, T. R., K. A. Haynes, and R. A. Barnes.** 1990. Value of antigen detection in predicting invasive aspergillosis. Lancet 336:1210-1213.
33. **Sandhu, G. S., B. C. Kline, L. Stockman, and G. D. Roberts.** 1995. Molecular probes for diagnosis of fungal infections. J. Clin. Microbiol. 33:2913-2919.
34. **Shin, J. H., F. S. Nolte, and C. J. Morrison.** 1997. Rapid identification of *Candida* species in blood cultures using a clinically useful PCR method. J. Clin. Microbiol. *in press*.
35. **Tang, C. M., D. W. Holden, A. Aufauvre-Brown, and J. Cohen.** The detection of *Aspergillus* spp. by the polymerase chain reaction and its evaluation in bronchoalveolar lavage fluid. Amer. Rev. Respir. Dis 148:1313-1317.

36. **Thompson, B. H., W. Stanford, J. R. Galvin, and Y. Kurlhara.** 1995. Varied radiologic appearances of pulmonary aspergillosis. *Radiographics* 15:1273-1284.
37. **Tierney, Jr. L.M.** Aspergillosis. *In* Current Medical Diagnosis and Treatment. 33rd ed. Norwalk, Conn.: Appleton and Lange, 1994.
38. **Verweij, P. E., J. P. Latge, A. J. Rijs, W. J. Melchers, B. E. De Pauw, J. A. Hoogkamp-Korstanje, and J. F. Mels.** 1995. Comparison of antigen detection and PCR assay using bronchoalveolar lavage fluid for diagnosing invasive pulmonary aspergillosis in patients receiving treatment for hematological malignancies. *J. Clin. Microbiol.* 33:3150-3153.
39. **von Eiff, M., N. Roos, R. Schulten, M. Hesse, M. Zuhisdorf, and J. van de Loo.** 1995. Pulmonary aspergillosis: early diagnosis improves survival. *Respiration* 62:341-347.
40. **von Eiff, M., N. Roos, W. Fegeler, C. von Eiff, R. Schulten, M. Hesse, M. Zuhisdorf, and J. van de Loo.** 1996. Hospital acquired *Candida* and *Aspergillus* pneumonia - diagnostic approaches and clinical findings. *J. Hosp. Infect.* 32:17-28.
41. **Walsh, T. J.** 1993. Management of immunocompromised patients with evidence of an invasive mycosis. *Hemat. Oncol. Clin. N .Amer.* 7:1003-1026.
42. **Walsh, T. J., C. Gonzalez, C. A. Lyman, S. J. Chanock, and P. A. Pizzo.** 1996. Invasive fungal infections in children: recent advances in diagnosis and treatment. *Adv. Ped. Inf. Dis.* 11:187-290.
43. **Walsh, T. J., B. De Pauw, E. Anaissle, and P. Martino.** 1994. Recent advances in the epidemiology, prevention, and treatment of invasive fungal infections in neutropenic patients. *J. Med. Vet. Mycol.* 32 Supp 1:33-51.
44. **Warnock, D. W.** 1995. Fungal complications of transplantation: diagnosis, treatment, and prevention. *J. Antimicrob. Chemother.* 36 Suppl B:73-90.

45. **Yamakami, Y., A. Hashimoto, I. Tokimatsu, and M. Nasu.** 1996. PCR detection of DNA specific for *Aspergillus* species in serum of patients with invasive aspergillosis. *J. Clin. Microbiol.* **34**:2464-2468.
- 5 46. **Young, R. C., and J. E. Bennett.** 1971. Invasive aspergillosis: absence of detectable antibody response. *Am. Rev. Respir. Dis* **104**:710-716.
- 10 47. **Zervos, M. J. and J. A. Vasquez.** 1996. DNA analysis in the study of fungal infections in the immunocompromised host. *Clin. Lab. Med.* **16**:73-88.

CLAIMS

We claim:

1. An isolated nucleic acid probe for identifying a species selected from the group consisting of *Aspergillus flavus* (SEQ ID NO:1), *Aspergillus fumigatus* (SEQ ID NO:2), *Aspergillus niger* (SEQ ID NO:3), *Aspergillus terreus* (SEQ ID NO:4), *Aspergillus nidulans* (SEQ ID NO:5), *Fusarium solani* (SEQ ID NO:6), *Fusarium moniliforme* (SEQ ID NO:7), *Mucor rouxii* (SEQ ID NO:8), *Mucor racemosus* (SEQ ID NO:9), *Mucor plumbeus* (SEQ ID NO:10), *Mucor indicus* (SEQ ID NO:11), *Mucor circinilloides f. circinelloides* (SEQ ID NO:12), *Rhizopus oryzae* (SEQ ID NO:13 and NO:14), *Rhizopus microsporus* (SEQ ID NO:15 and 16), *Rhizopus circinans* (SEQ ID NO:17 and 18), *Rhizopus stolonifer* (SEQ ID NO: 19), *Rhizomucor pusillus* (SEQ ID NO:20), *Absidia corymbifera* (SEQ ID NO:21 and 22), *Cunninghamella elegans* (SEQ ID NO:23), *Pseudallescheria boydii* (teleomorph of *Scedosporium apiospermum*) (SEQ ID NO:24, 25, 26, and 27), *Penicillium notatum* (SEQ ID NO:28), or *Sporothrix schenckii* (SEQ ID NO:29) wherein the probe selectively hybridizes to a portion of the nucleic acid of SEQ ID NOS:1-29, or a complementary sequence thereof, respectively.

2. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with an *Aspergillus flavus* nucleic acid of SEQ ID NO:1, or a complementary sequence thereof.

3. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with an *Aspergillus fumigatus* nucleic acid of SEQ ID NO:2, or a complementary sequence thereof.

4. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with an *Aspergillus niger* nucleic acid of SEQ ID NO:3, or a complementary sequence thereof.

5. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with an *Aspergillus terreus* nucleic acid of SEQ ID NO:4, or a complementary sequence thereof.

6. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with an *Aspergillus nidulans* nucleic acid of SEQ ID NO:5, or a complementary sequence thereof.

7. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Fusarium solani* nucleic acid of SEQ ID NO:6, or a complementary sequence thereof.

8. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Fusarium moniliforme* of SEQ ID NO:7, or a complementary sequence thereof.

9. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Mucor rouxii* of SEQ ID NO:8, or a complementary sequence thereof.

10. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Mucor racemosus* of SEQ ID NO:9, or a complementary sequence thereof.

11. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Mucor plumbeus* of SEQ ID NO:10, or a complementary sequence thereof.

12. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Mucor indicus* of SEQ ID NO:11, or a complementary sequence thereof.

13. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Mucor circinilloides f. circinelloides* of SEQ ID NO:12, or a complementary sequence thereof.

14. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Rhizopus oryzae* of SEQ ID NO:13 and 14, or a complementary sequence thereof.

15. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Rhizopus microsporus* of SEQ ID NO:15 and 16, or a complementary sequence thereof.

16. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Rhizopus circinans* of SEQ ID NO:17 and 18, or a complementary sequence thereof.

17. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Rhizopus stolonifer* of SEQ ID NO:19, or a complementary sequence thereof.

18. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Rhizomucor pusillus* of SEQ ID NO:20, or a complementary sequence thereof.

19. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Absidia corymbifera* of SEQ ID NO:21 and 22, or a complementary sequence thereof.

20. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Cunninghamella elegans* of SEQ ID NO:23, or a complementary sequence thereof.

21. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Pseudallescheria boydii* (teleomorph of *Scedosporium apiospermum*) of SEQ ID NO:24, 25, 26 and 27, or a complementary sequence thereof.

22. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Penicillium notatum* of SEQ ID NO:28, or a complementary sequence thereof.

23. The isolated nucleic acid probe of Claim 1 capable of selectively hybridizing with a *Sporothrix schenkii* of SEQ ID NO:29, or a complementary sequence thereof.

24. A method of detecting a species selected from the group consisting of *Aspergillus flavus* (SEQ ID NO:1), *Aspergillus fumigatus* (SEQ ID NO:2), *Aspergillus niger* (SEQ ID NO:3), *Aspergillus terreus* (SEQ ID NO:4), *Aspergillus nidulans* (SEQ ID NO:5), *Fusarium solani* (SEQ ID NO:6), *Fusarium moniliforme* (SEQ ID NO:7), *Mucor rouxii* (SEQ ID NO:8), *Mucor racemosus* (SEQ ID NO:9), *Mucor plumbeus* (SEQ ID NO:10), *Mucor indicus* (SEQ ID NO:11), *Mucor circinilloides f. circinelloides* (SEQ ID NO:12), *Rhizopus oryzae* (SEQ ID NO:13 and NO:14), *Rhizopus microsporus* (SEQ ID NO:15 and 16), *Rhizopus circinans* (SEQ ID NO:17 and 18), *Rhizopus stolonifer* (SEQ ID NO: 19), *Rhizomucor pusillus* (SEQ ID NO:20), *Absidia corymbifera* (SEQ ID NO:21 and 22), *Cunninghamella elegans* (SEQ ID NO:23), *Pseudallescheria boydii* (teleomorph of *Scedosporium apiospermum*) (SEQ ID NO:24, 25, 26, and 27), *Penicillium notatum* (SEQ ID NO:28), or *Sporothrix schenkii* (SEQ ID NO:29) in a sample comprising combining the sample with a nucleic acid probe capable of selectively hybridizing with a nucleic acid of SEQ ID NO:1-29, or a complementary sequence thereof, respectively, the presence of hybridization indicating the detection of the species in the sample.

25. The method of Claim 24, wherein the probe is capable of selectively hybridizing with an *Aspergillus flavus* nucleic acid of SEQ ID NO:1, or a complementary sequence thereof.

26. The method of Claim 24, wherein the probe is capable of selectively hybridizing with an *Aspergillus fumigatus* nucleic acid of SEQ ID NO:2, or a complementary sequence thereof.

27. The method of Claim 24, wherein the probe is capable of selectively hybridizing with an *Aspergillus niger* nucleic acid of SEQ ID NO:3, or a complementary sequence thereof.

28. The method of Claim 24, wherein the probe is capable of selectively hybridizing with an *Aspergillus terreus* nucleic acid of SEQ ID NO:4, or a complementary sequence thereof.

29. The method of Claim 24, wherein the probe is capable of selectively hybridizing with an *Aspergillus nidulans* nucleic acid of SEQ ID NO:5, or a complementary sequence thereof.

30. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Fusarium solani* nucleic acid of SEQ ID NO:6, or a complementary sequence thereof.

31. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Fusarium moniliforme* of SEQ ID NO:7, or a complementary sequence thereof.

32. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Mucor rouxii* of SEQ ID NO:8, or a complementary sequence thereof.

33. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Mucor racemosus* of SEQ ID NO:9, or a complementary sequence thereof.

34. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Mucor plumbeus* of SEQ ID NO:10, or a complementary sequence thereof.

35. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Mucor indicus* of SEQ ID NO:11, or a complementary sequence thereof.

36. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Mucor circinilloides* f. *circinelloides* of SEQ ID NO:12, or a complementary sequence thereof.

37. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Rhizopus oryzae* of SEQ ID NO:13 and 14, or a complementary sequence thereof.

38. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Rhizopus microsporus* of SEQ ID NO:15 and 16, or a complementary sequence thereof.

39. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Rhizopus circinans* of SEQ ID NO:17 and 18, or a complementary sequence thereof.

40. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Rhizopus stolonifer* of SEQ ID NO:19, or a complementary sequence thereof.

41. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Rhizomucor pusillus* of SEQ ID NO:20, or a complementary sequence thereof.

42. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Absidia corymbifera* of SEQ ID NO:21 and 22, or a complementary sequence thereof.

43. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Cunninghamella elegans* of SEQ ID NO:23, or a complementary sequence thereof.

44. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Pseudallescheria boydii* (teleomorph of

Scedosporium apiospermum) of SEQ ID NO:24, 25, 26 and 27, or a complementary sequence thereof.

45. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Penicillium notatum* of SEQ ID NO:28, or a complementary sequence thereof.

46. The method of Claim 24, wherein the probe is capable of selectively hybridizing with a *Sporothrix schenkii* of SEQ ID NO:29, or a complementary sequence thereof.

47. An isolated nucleic acid probe for identifying a member of a genus selected from the group consisting of *Aspergillus*, *Fusarium* and *Mucor* wherein the probe selectively hybridizes to a portion of the nucleic acid of SEQ ID NOS:58-60, or a complementary sequence thereof, respectively.

48. An isolated nucleic acid probe for identifying a fungus wherein the probe selectively hybridizes to a portion of the nucleic acid of SEQ ID NO:61, or a complementary sequence thereof, respectively.

49. A method for detecting a member of a genus selected from the group consisting of *Aspergillus*, *Fusarium* and *Mucor* in a sample comprising combining the sample with a nucleic acid probe capable of selectively hybridizing to a portion of the nucleic acid of SEQ ID NOS:58-60, or a complementary sequence thereof, respectively, the presence of hybridization indicating the detection of the respective genus.

50. A method for detecting a fungus in a sample comprising combining the sample with a nucleic acid probe capable of selectively hybridizing to a portion of the nucleic acid of SEQ ID NO:61, or a complementary sequence thereof, respectively, the presence of hybridization indicating the detection of the fungus.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.